

AD-A048 973

CALIFORNIA UNIV LOS ANGELES DEPT OF CHEMISTRY  
SYNTHESIS OF ELECTRON-DEFICIENT BIFFERRACARBORANES. (U)  
JAN 78 C G SALENTINE, M F HAWTHORNE

F/G 7/3

UNCLASSIFIED

TR-98

N00014-76-C-0390

NL

| 0 |

ADAO48 973



END  
DATE  
FILMED

2 -78

DDC

AD A048973

AD No.             
DDC FILE COPY

OFFICE OF NAVAL RESEARCH

Contract No. 15 N00014-76-C-0390

Task No. NR 053-608

9 TECHNICAL REPORT, NO. 98

14 TR-98

6 Synthesis of Electron-Deficient Biferracarboranes.

By

10 C. G. Salentine and M. F. Hawthorne

Prepared for Publication  
in  
Inorganic Chemistry

Department of Chemistry  
University of California  
Los Angeles, California 90024

11 10 January 20, 1978

DDC  
RECEIVED  
JAN 20 1978  
F

Reproduction in whole or part is permitted for  
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited

1473

072 255

4B

[Contribution No. 3829 from the Department of Chemistry,  
University of California, Los Angeles, California 90024

Synthesis of Electron-Deficient Biferracarboranes

By

C. G. Salentine and M. F. Hawthorne\*

Department of Chemistry  
University of California  
Los Angeles, California 90024

### Abstract

The polyhedral expansion of 3,1,2- $C_5H_5FeC_2B_9H_{11}$  produced three new electron-deficient biferracarboranes  $(C_5H_5Fe)_2C_2B_9H_{11}$ ,  $(C_5H_5Fe)_2C_2B_8H_9(OH)$  and  $[C_5H_5FeC_2B_9H_{11}FeC_2B_9H_{11}]^-$ . All three complexes contained formal Fe(III) atoms and were diamagnetic. The proposed structures, based on spectroscopic and crystallographic data, contain non-bonded iron atoms. These complexes constitute the first examples of long-range electron-spin coupling through a carborane polyhedron.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTI DATE	
BY	
DISSEMINATION/AVAILABILITY CODES	
SPECIAL	
A	



Ferracarboranes have been synthesized containing formal Fe(II) and Fe(III) oxidation states.<sup>1</sup> Those of formula  $[\text{Fe}(\text{C}_2\text{B}_n\text{H}_{n+2})_2]^{2-}$  and  $[\text{C}_5\text{H}_5\text{FeC}_2\text{B}_n\text{H}_{n+2}]^-$  contain formal Fe(II) and are readily air-oxidized to the stable Fe(III) complexes  $[\text{Fe}(\text{C}_2\text{B}_n\text{H}_{n+2})_2]^-$  and  $\text{C}_5\text{H}_5\text{FeC}_2\text{B}_n\text{H}_{n+2}$ . These latter Fe(III) metallocarboranes can be viewed as two-carbon carboranes in which a  $\{\text{C}_5\text{H}_5\text{Fe}\}$  vertex has replaced a  $\{\text{BH}\}$  vertex. Because  $\{\text{C}_5\text{H}_5\text{Fe(II)}\}$  is formally "isoelectronic" with  $\{\text{BH}\}$ , the Fe(III) complexes represent polyhedra containing one electron less than the required  $2n+2$  framework electrons.<sup>2</sup> A result of this electron deficiency is that polyhedral rearrangements have not yet been observed in this class of metallocarboranes. Crystallographic studies<sup>1d</sup> have shown that these Fe(III) metallocarboranes possess regular closo-polyhedral geometries, indicating that one-electron deficiency has little or no effect upon polyhedral structure.

Prior to this work, only one electron-deficient biferracarborane had been reported,<sup>3</sup> the two-electron deficient species  $(\text{C}_5\text{H}_5\text{Fe}^{\text{III}})_2\text{C}_2\text{B}_6\text{H}_8$  prepared by polyhedral expansion of 4,5- $\text{C}_2\text{B}_7\text{H}_9$ . This curious compound was initially obtained in a paramagnetic ( $\mu_{\text{eff}} = 3.05$  B.M.) form presumably containing two unpaired electrons. It underwent a slow transformation to a diamagnetic species and a crystal structure<sup>3</sup> of the diamagnetic complex showed a unique 10-vertex closo-polyhedral geometry incorporating an iron-iron bond ( $2.571(1) \text{ \AA}$ ), much different from the bicapped square antiprismatic geometry usually observed in 10-vertex carboranes and metallocarboranes. The structure was described as a derivation of a tri-capped trigonal prism, the next lower polyhedron for which the biferracarborane did satisfy the electronic requirements. Recently, structurally novel 14-vertex nido-biferracarboranes which do satisfy the polyhedral electronic requirements were reported.<sup>4</sup>


As a continuation of our studies<sup>5</sup> of electronic effects upon geometry in metallocarborane polyhedra, we report here the synthesis of several new electron-

deficient biferracarboranes incorporating two formal iron (III) metal atoms into twelve- and thirteen-vertex metallocarborane polyhedra.


### Results and Discussion

#### The Polyhedral Expansion of 3-( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)-3-Fe-1,2-C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>.

The reduction of 3-( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)-3-Fe-1,2-C<sub>2</sub>B<sub>9</sub>H<sub>11</sub> with three equivalents of sodium followed by treatment with NaC<sub>5</sub>H<sub>5</sub> and FeCl<sub>2</sub> afforded a mixture of products. Two neutral biferracarboranes were isolated by column chromatography on silica gel, and a third anionic bimetallic species was precipitated from water as the tetramethylammonium salt.



I



II

III

The golden orange-brown crystalline I exhibited a mass spectrum with a cut-off at *m/e* 376 corresponding to the <sup>12</sup>C<sub>12</sub><sup>1</sup>H<sub>21</sub><sup>11</sup>B<sub>9</sub><sup>56</sup>Fe<sub>2</sub><sup>+</sup> ion. Elemental analysis also confirmed this empirical formula. The diamagnetism of I was indicated from its nmr spectra. The 60 MHz <sup>1</sup>H nmr spectrum (in CD<sub>2</sub>Cl<sub>2</sub>) showed a sharp resonance at  $\tau$  5.14 and a broad resonance at  $\tau$  10.81 of relative areas 5:1. These were assigned to equivalent cyclopentadienyl groups and equivalent carborane C-H groups, respectively. The 80.5 MHz <sup>11</sup>B nmr spectrum contained resonances of area 1:2:4:2 at -113.1, -42.4, -22.4 and -5.7 ppm, relative to Et<sub>2</sub>O·BF<sub>3</sub>. This high degree of symmetry within a 13-vertex polyhedron considerably limits the structural possibilities. Although fluxionality within a less symmetric polyhedron<sup>1c</sup> is one possible explanation of the nmr data, the <sup>11</sup>B nmr spectrum of I was unchanged to -75°. The resonance at -113 ppm almost certainly arises from a low-coordinate

boron atom adjacent to two metal vertices (see references 3 and 5a). The proposed fluxional process<sup>1C</sup> would "average" the low-coordinate boron position in the intermediate structure and a very low field resonance would not be expected in the <sup>11</sup>B nmr spectrum. Based on the above, we therefore postulate that I is not fluxional in solution, in contrast to other 13-vertex metallocarboranes such as 4-( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)-4-Co-1,6-C<sub>2</sub>B<sub>10</sub>H<sub>12</sub><sup>1C</sup> and 4,5-( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>-4,5-Co<sub>2</sub>-1,6-C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>.<sup>6</sup>

The observed diamagnetism of I would tend to imply the existence of a metal-metal bond;<sup>3</sup> however, preliminary X-ray crystallographic results<sup>7</sup> indicate an Fe-Fe distance of  $\sim 3.20 \text{ \AA}$ . The preceding considerations lead to a proposed structure for I, 4,5-( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>-4,5-Fe<sub>2</sub>-2,3-C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>, as shown in Figure 1. The metal atoms reside at their favored high-coordinate positions

---

Figure 1

---

in the 13-vertex polyhedron. The structure with carbon atoms at the 10,11 positions is not possible because replacement of one C<sub>5</sub>H<sub>5</sub><sup>-</sup> ligand with a C<sub>2</sub>B<sub>9</sub>H<sub>11</sub><sup>2-</sup> ligand (complex II, *vide infra*) would render the carborane C-H groups nonequivalent in II, and they remain equivalent. The 12,13 positions for the carbon atoms are less likely because the starting material contained both carbon atoms adjacent to iron.

The low-coordinate 1-position in the structure proposed for I is occupied by boron - this is the first 13-vertex closo-metallocarborane synthesized that does not have carbon at its favored low-coordinate position.<sup>8</sup> Isolation of such a thermodynamically unstable species is possible with the electron-deficient ferracarboranes because they have been found not to undergo polyhedral rearrangement.



rearrangement.

The proposed geometry for the two electron-deficient I has the same gross structure as found for  $C_5H_5CoC_2B_{10}H_{12}$  which obeys the  $2n+2$  electron rule. Considering the structural result of  $(C_5H_5Fe)_2C_2B_6H_8$ ,<sup>3</sup> it is possible that I could adopt a different polyhedral geometry. While this proposed structure is necessarily tentative, it agrees well with the available data. A single-crystal X-ray diffraction study of complex II has been initiated and may provide further structural information.

Complex I represents the first diamagnetic metallocarborane containing two paramagnetic centers not within bonding distance. The diamagnetism appears to be a result of spin pairing by interactions of the two iron atoms through the carborane ligand. Studies by Taube<sup>9</sup> and Day<sup>10</sup> have suggested that two metals can interact strongly through conjugated pi-electron systems. The first example of complete electron delocalization through a bridging ligand in an organometallic complex was shown<sup>11</sup> recently for the bisfulvalenediiron dication,  $[(C_{10}H_8)_2Fe_2]^{2+}$ , which contains two Fe(III) centers separated by about 3.9 Å, and yet is diamagnetic. The cyclic voltammogram of I showed a reversible reduction at  $E_{p/2} = -0.59V$  and an irreversible oxidation at  $E_{p/2} = +1.36V$ . The reduction is presumably to a mixed-valence Fe(II)-Fe(III) system. Further investigation of the reduced species is under way.


The second product isolated from the polyhedral expansion reaction was the diamagnetic green complex, II. Elemental analysis was consistent with the formulation  $[(CH_3)_4N][C_5H_5FeC_2B_9H_{11}FeC_2B_9H_{11}]$ . The 60 MHz  $^1H$  nmr spectrum, measured in  $d_6$ -acetone, exhibited resonances of area 12:5:2:2 at  $\tau$  6.57,  $\tau$  5.06,  $\tau$  7.50 and  $\tau$  10.27, assigned to tetramethylammonium, cyclopentadienyl, carborane C-H and carborane C-H protons, respectively. The carborane C-H resonance at



$\tau$  7.50 was within about 1 ppm to that observed<sup>1a</sup> for  $[3-(\eta^5\text{-C}_5\text{H}_5)\text{-3-Fe-1,2-C}_2\text{B}_9\text{H}_{11}]^-$ , and the resonance at  $\tau$  10.27 was within 1/2 ppm to that observed for I. The 80.5 MHz  $^{11}\text{B}$  nmr spectrum of II also closely corresponded to a superposition of the  $^{11}\text{B}$  spectra of I and  $[3-(\eta^5\text{C}_5\text{H}_5)\text{-3-Fe-1,2-C}_2\text{B}_9\text{H}_{11}]^-$ . Doublets corresponding to the  $\eta^5\text{-1,2-C}_2\text{B}_9\text{H}_{11}^{2-}$  ligand were observed in areas 1:1:4:2:1 at -7.7, +0.5, +4.8, +17.5 and +20.6 ppm. Resonances corresponding to the  $\text{C}_2\text{B}_9\text{H}_{11}^{4-}$  bimetallic cage were observed in areas 1:1:1:2:2:2 at -110.1, -46.1, -39.8, -23.9, -22.6 and -5.8 ppm, assigned by comparison with the  $^{11}\text{B}$  nmr spectrum of I. The area 4 resonance in the  $^{11}\text{B}$  nmr spectrum of I, at -22.4 ppm, was split into two area 2 peaks in II at -23.9 and -22.6 ppm. Likewise, the area 2 resonance of I at -42.4 ppm was split into two area 1 peaks in II at -46.1 and -39.8 ppm. These data are consistent with a structure for II entirely analogous to I, with a  $\text{C}_5\text{H}_5^-$  ligand replaced by a  $1,2\text{-C}_2\text{B}_9\text{H}_{11}^{2-}$  ligand (see Figure 1). This substitution allows an assignment of the  $^{11}\text{B}$  nmr spectrum of complex I: the area 1 resonance at -113.1 ppm is due to B1; the area 2 resonance at -42.4 ppm is due to B10, B11; the area 4 resonance at -22.4 ppm is due to B6, B7, B8, B9; and the area 2 resonance at -5.7 ppm is due to B12, B13. Thus II is formulated as  $[(\text{CH}_3)_4\text{N}][4-(\eta^5\text{-C}_5\text{H}_5)\text{-5-(}\eta^5\text{-1,2-C}_2\text{B}_9\text{H}_{11})\text{-4,5-Fe}_2\text{-2,3-C}_2\text{B}_9\text{H}_{11}]$ .

The third product of the reaction was a brown crystalline material, isolated in low yield by column chromatography,  $(\text{C}_5\text{H}_5\text{Fe})_2\text{C}_2\text{B}_8\text{H}_9(\text{OH})$ , III. The mass spectrum showed a cutoff at  $m/e$  380 corresponding to the  $^{12}\text{C}_{12}^{1}\text{H}_{20}^{11}\text{B}_8^{56}\text{Fe}_2^{16}\text{O}^+$  ion. A high resolution mass spectrum confirmed the formula; calcd for  $^{12}\text{C}_{11}^{13}\text{C}^1\text{H}_{20}^{11}\text{B}_8^{57}\text{Fe}_2^{16}\text{O}^+$ : 383.1000, found: 383.1003. This biferracarborane was also diamagnetic, showing one cyclopentadienyl resonance in the  $^1\text{H}$  nmr spectrum ( $d_6$ -acetone) at  $\tau$  4.99. The 80.5 MHz  $^{11}\text{B}$  nmr spectrum contained doublets of area 2:2:2:1:1 at -26.4, -19.3, -17.3, -3.6 and -0.8 ppm and indicated that OH substitution had occurred at a carborane carbon atom. The infrared spectrum contained a peak at  $\sim 3500\text{ cm}^{-1}$  assigned to  $\nu_{\text{O-H}}$ . Assuming a gross icosahedral

geometry for III, there are only four possibilities for the structure:



The first two possibilities contain adjacent iron atoms, while the last two contain adjacent carbon atoms. Because the starting material contained adjacent carbon atoms, and it has been shown (*vide supra*) that a direct iron-iron bond is not necessary for diamagnetism, we favor the last two possibilities over the first two. Also, because the carbon atoms were adjacent to iron in the starting material,  $1\text{-OH-}3,6\text{-(}\eta^5\text{-C}_5\text{H}_5\text{)}_2\text{-}3,6\text{-Fe}_2\text{-}1,2\text{-C}_2\text{B}_8\text{H}_9$  is the most probable structure of III. Traces of water or hydroxide in the polyhedral expansion reaction mixture could have resulted in the production of III, analogous to the synthesis of  $3,6\text{-(}\eta^5\text{-C}_5\text{H}_5\text{)}_2\text{-}3,6\text{-Co}_2\text{-}1,2\text{-C}_2\text{B}_8\text{H}_{10}$  by treatment of  $3\text{-(}\eta^5\text{-C}_5\text{H}_5\text{)-}3\text{-Co-}1,2\text{-C}_2\text{B}_9\text{H}_{11}$  with KOH,  $\text{C}_5\text{H}_5^-$  and  $\text{CoCl}_2$ .<sup>12</sup> The geometry found for III could also be easily generated from I by removal of the low-coordinate boron position with the formation of the C2-C3 bond (Figure 1). As with I, it is possible that the geometry of III may deviate significantly from icosahedral, as III is also two-electron deficient.

Isolation of a stable derivative of  $(\text{C}_5\text{H}_5\text{Fe})_2\text{C}_2\text{B}_8\text{H}_{10}$  led to attempts to prepare the parent compound. The polyhedral expansion of  $1\text{-(}\eta^5\text{-C}_5\text{H}_5\text{)-}1\text{-Fe-}2,3\text{-C}_2\text{B}_8\text{H}_{10}$  with  $\text{C}_5\text{H}_5^-$  and  $\text{FeCl}_2$  produced unstable ferracarboranes which rapidly decomposed to ferrocene in solution; use of  $[\text{C}_5\text{H}_5\text{FeC}_6\text{H}_6]^+$  as a  $\text{FeC}_5\text{H}_5$  source produced a red compound which eluted from silica gel in dichloromethane. The mass spectrum showed a cutoff at  $m/e$  364 corresponding to  $(\text{C}_5\text{H}_5\text{Fe})_2\text{C}_2\text{B}_8\text{H}_{10}$ . The 60 MHz  $^1\text{H}$  nmr spectrum contained two broad peaks of equal area in the cyclopentadienyl region and indicated that the compound was paramagnetic. Cyclic

voltammetry showed a reversible reduction at  $E_{p/2} = -0.56V$  and a quasi-reversible reduction at  $E_{p/2} = -1.40V$ . This complex was also unstable, even in the solid state, and decomposed in a few days to  $1-(\eta^5-C_5H_5)-1-Fe-2,3-C_2B_8H_{10}$ . Apparently, the bimetallocarborane resulting from polyhedral expansion of  $1-(\eta^5-C_5H_5)-1-Fe-2,3-C_2B_8H_{10}$ , which contains non-adjacent carbon atoms, did not have a stable isomeric arrangement of heteroatoms within the polyhedron. It is interesting that the only other paramagnetic biferracarborane,  $(C_6H_5Fe)_2C_2B_6H_8$ , was also unstable with respect to transformation to a diamagnetic form.

Among the results of these studies is the important observation that paramagnetic metal centers not within bonding distance may participate in long-range coupling through a carborane cage. The greater stability and ease of purification of complex I as compared to the bisfulvalendiiron dication<sup>11</sup> make it an attractive candidate for further physical studies.

## Experimental Section

### Physical Measurements

Ultraviolet-visible spectra were measured with a Cary 14 spectrophotometer. Infrared spectra were determined using a Perkin Elmer Model 137 sodium chloride spectrophotometer. Proton nmr spectra were obtained on a Varian A-60D spectrometer. The 80.5 MHz <sup>11</sup>B nmr spectra were obtained with an instrument designed by Professor F. A. L. Anet of this department. Electrochemical data were obtained on an instrument described previously.<sup>13</sup> Mass spectra were measured using an Associated Electrical Industries MS-9 spectrometer.

Elemental analyses were carried out by Schwarzkopf Microanalytical Laboratories, Woodside, N. Y.



## Materials

Literature methods were used to prepare  $3-(\eta^5\text{-C}_5\text{H}_5)\text{-3-Fe-1,2-C}_2\text{B}_9\text{H}_{11}$ , <sup>1a</sup>  
 $1-(\eta^5\text{-C}_5\text{H}_5)\text{-1-Fe-2,3-C}_2\text{B}_8\text{H}_{10}$  <sup>1b</sup> and  $[\text{C}_5\text{H}_5\text{FeC}_6\text{H}_6][\text{PF}_6]$ . <sup>14</sup> Anhydrous sublimed  
 ferric chloride and hydrogen reduced iron powder were obtained from Matheson,  
 Coleman and Bell. Tetrahydrofuran (THF) was distilled from lithium aluminum  
 hydride and stored under nitrogen prior to use. All other solvents were reagent  
 grade and used without further purification. Naphthalene, Spectroquality aceto-  
 nitrile and tetramethylammonium chloride were obtained from Matheson, Coleman  
 and Bell. Sodium hydride, as a 50% dispersion in mineral oil was obtained from  
 ROC/RIC Chemical Corp. Dicyclopentadiene was obtained from Aldrich Chemical Co.  
 and converted to cyclopentadiene immediately prior to use. THF solutions of  
 sodium cyclopentadienide were prepared as previously described <sup>1c</sup> and immediately  
 used. Sodium metal was purchased from Allied Chemical Co. Silica gel powder,  
 60-200 mesh, was obtained from J. T. Baker Chemical Co. for use in column chroma-  
 tography. Preparative thick-layer chromatography was performed with Chrom AR  
 Sheet 1000 purchased from Mallinckrodt Chemical Co.

## The Polyhedral Expansion of $3-(\eta^5\text{-C}_5\text{H}_5)\text{-3-Fe-1,2-C}_2\text{B}_9\text{H}_{11}$

Into a nitrogen-flushed three-necked 500 ml round-bottomed flask equipped  
 with a mechanical stirrer and nitrogen inlet was placed 2.53 g (10.0 mmol)  
 $3,1,2\text{-C}_5\text{H}_5\text{FeC}_2\text{B}_9\text{H}_{11}$  followed by 100 ml THF. To this was added freshly cut  
 sodium metal (0.70 g, 30 mmol) and 1.0 g naphthalene. The solution was stirred  
 six days under nitrogen, yielding an orange-green color with all the sodium con-  
 sumed. To this solution was added a THF solution of 35 mmol  $\text{NaC}_5\text{H}_5$  followed by  
 a THF slurry of 30 mmol  $\text{FeCl}_2$  (prepared by refluxing 0.70 g Fe powder with 3.24 g  
 $\text{FeCl}_3$  for 18 hr under nitrogen). The mixture was stirred 16 hr under nitrogen,  
 then oxygen was bubbled through the solution for 20 min. The solution was fil-  
 tered through Celite and added to 40 g silica gel. Solvent was removed and the



solids were placed atop a 4 x 30 cm silica gel chromatography column prepared in hexane. Elution with hexane produced a yellow band containing 1.6 g ferrocene. Elution with 20% CH<sub>2</sub>Cl<sub>2</sub>/hexane developed a large red-purple band which yielded 0.98 g of a mixture of starting material and (C<sub>5</sub>H<sub>5</sub>Fe)<sub>2</sub>C<sub>2</sub>B<sub>9</sub>H<sub>11</sub>, I. These were separated by thick-layer chromatography to yield 290 mg of the less polar starting material and 647 mg (17%) of I as shiny metallic copper-colored plates, dec. above ~180° without melting. Anal. Calcd for C<sub>12</sub>H<sub>21</sub>B<sub>9</sub>Fe<sub>2</sub>: C, 38.51; H, 5.66; B, 26.00; Fe, 29.84. Found: C, 38.15; H, 5.50; B, 25.00; Fe, 28.61. High resolution mass measurement; calcd for <sup>12</sup>C<sub>12</sub><sup>1</sup>H<sub>21</sub><sup>11</sup>B<sub>9</sub><sup>56</sup>Fe<sub>2</sub><sup>+</sup>: 376.1168. Found: 376.1202. Infrared spectrum (Nujol mull): 3070(w), 2600(w), 2510(vs), 2470(vs), 1420(s), 1120(w), 1130(w), 1070(m), 1060(w) 1025(s), 1010(m), 965(m), 935(w), 925(w), 910(w), 905(w), 885(m), 875(w), 860(w), 850(s), 835(s), 755(w), 738(m), 728(m), 720(m) cm<sup>-1</sup>.

A yellow-brown band was eluted from the column in 60% CH<sub>2</sub>Cl<sub>2</sub>/hexane which upon rotary evaporation of solvent produced 60 mg (1.6%) of green-brown crystals of III, (C<sub>5</sub>H<sub>5</sub>Fe)<sub>2</sub>C<sub>2</sub>B<sub>8</sub>H<sub>9</sub>OH, m.p. >300°. Anal. Calcd for <sup>12</sup>C<sub>12</sub><sup>1</sup>H<sub>20</sub><sup>11</sup>B<sub>8</sub><sup>56</sup>Fe<sub>2</sub><sup>16</sup>O<sup>+</sup>: 380.0957. Found: 380.0973. Calcd for <sup>12</sup>C<sub>11</sub><sup>13</sup>C<sup>1</sup>H<sub>20</sub><sup>11</sup>B<sub>8</sub><sup>57</sup>Fe<sub>2</sub><sup>16</sup>O<sup>+</sup>: 383.1000. Found: 383.1003. Infrared spectrum (Nujol mull): 3450(w), 3090(w), 2490(vs), 1425(m), 1415(m), 1275(m), 1150(m), 1110(s), 1070(s), 1045(w), 1020(m), 1015(m), 988(m), 938(w), 882(m), 844(s), 720(w), 688(m) cm<sup>-1</sup>.

The silica gel was then extracted with CH<sub>3</sub>CN, yielding a deep green solution. Removal of CH<sub>3</sub>CN followed by addition of H<sub>2</sub>O, filtration and addition of excess (CH<sub>3</sub>)<sub>4</sub>NCl produced a green precipitate. The precipitate (150 mg) was dissolved in acetone and stripped onto silica gel and chromatographed on a 2 x 15 cm silica gel column. Elution with CH<sub>2</sub>Cl<sub>2</sub> removed impurities while elution with 10% acetone/CH<sub>2</sub>Cl<sub>2</sub> eluted the green band. Addition of *n*-propanol to the filtrate followed by slow rotary evaporation produced 84 mg (1.6%) of dark green crystals of II, [(CH<sub>3</sub>)<sub>4</sub>N][C<sub>5</sub>H<sub>5</sub>FeC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>FeC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>], m.p. >300°. Anal. Calcd for C<sub>13</sub>H<sub>39</sub><sup>-</sup>B<sub>18</sub>Fe<sub>2</sub>N: C, 30.28; H, 7.62; B, 37.73; Fe, 21.66; N, 2.72. Found: C, 30.17;

H, 7.76; B, 36.51; Fe, 21.74; N, 2.69. Infrared spectrum (Nujol mull): 3100(w), 2500(vs), 1490(s), 1425(m), 1290(w), 1155(w), 1125(w), 1070(m), 1060(w), 1035(m), 1025(m), 995(m), 977(m), 950(m), 883(m), 860(w), 840(s), 760(m), 740(m), 724(m)  $\text{cm}^{-1}$ .

Polyhedral Expansion of 1-( $\eta^5\text{-C}_5\text{H}_5$ )-1-Fe-2,3- $\text{C}_2\text{B}_8\text{H}_{10}$ .

This reaction was performed as above with 0.98 g (4.06 mmol) 1,2,3- $\text{C}_5\text{H}_5\text{-FeC}_2\text{B}_8\text{H}_{10}$ . To the reduced solution was added, as a solid, 6.5 g (19 mmol) [ $\text{C}_5\text{H}_5\text{FeC}_6\text{H}_6$ ][PF<sub>6</sub>]. The mixture was stirred 120 hr under nitrogen and worked up as above. Elution of the chromatography column with hexane produced ferrocene, and elution with 20%  $\text{CH}_2\text{Cl}_2$ -hexane yielded 60 mg starting material. A red band eluted in  $\text{CH}_2\text{Cl}_2$ . Addition of a small amount of hexane followed by rotary evaporation of the solvent produced 90 mg red-purple crystals, m.p.  $>310^\circ$ , whose mass and  $^1\text{H}$  nmr spectra were consistent with the formulation  $(\text{C}_5\text{H}_5\text{Fe})_2\text{C}_2\text{B}_8\text{H}_{10}$ . No 80.5 MHz  $^{11}\text{B}$  nmr spectrum was observable for the paramagnetic species. The elemental analysis and magnetic susceptibility were close to that expected for  $\text{C}_5\text{H}_5\text{FeC}_2\text{B}_8\text{H}_{10}$ , due presumably to solid state decomposition.

Acknowledgement

We are grateful to the Office of Naval Research for financial support.

### References

1. See, for example, the following: (a) M. F. Hawthorne, et al., J. Amer. Chem. Soc., 90, 879 (1968); (b) W. J. Evans, G. B. Dunks, and M. F. Hawthorne, ibid., 95, 4565 (1973); (c) D. F. Dustin, G. B. Dunks, and M. F. Hawthorne, ibid., 95, 1109 (1973); (d) A. Zalkin, D. H. Templeton, and T. E. Hopkins, ibid., 87, 3988 (1965).
2. R. W. Rudolph, Acc. Chem. Res., 9, 446 (1976).
3. K. P. Callahan, W. J. Evans, F. Y. Lo, C. E. Strouse, and M. F. Hawthorne, J. Amer. Chem. Soc., 97, 296 (1975).
4. (a) W. M. Maxwell, R. F. Bryan, and R. N. Grimes, J. Amer. Chem. Soc., 99, 4008 (1977); (b) W. M. Maxwell, R. Weiss, E. Sinn, and R. N. Grimes, ibid., 99, 4016 (1977).
5. (a) C. G. Salentine, C. E. Strouse, and M. F. Hawthorne, Inorg. Chem., 15, 1832 (1976); (b) C. G. Salentine and M. F. Hawthorne, ibid., 15, 2872 (1976).
6. W. J. Evans and M. F. Hawthorne, Inorg. Chem., 13, 869 (1974).
7. K. P. Callahan, preliminary crystallographic study of complex II.
8. M. F. Hawthorne, K. P. Callahan, and R. J. Wiersema, Tetrahedron, 30, 1795 (1974).
9. C. Creutz and H. Taube, J. Amer. Chem. Soc., 95, 1086 (1973), and references therein.
10. B. Mayoh and P. Day, J. Amer. Chem. Soc., 94, 2885 (1972).
11. C. LeVanda, K. Bechgaard, D. O. Cowan, U. T. Mueller-Westerhoff, P. Eilbracht, G. A. Candela, and R. L. Collins, J. Amer. Chem. Soc., 98, 3181 (1976).
12. J. Plešek, B. Štibr, and S. Heřmánek, Syn. Inorg. Metal. Chem., 3, 291 (1973).
13. T. E. Paxson, M. K. Kaloustian, G. M. Tom, R. J. Wiersema, and M. F. Hawthorne, J. Amer. Chem. Soc., 94, 4882 (1972).
14. R. B. King, "Organometallic Syntheses," Vol. 1, Academic Press, New York, N. Y., 1965, p. 138.



Table I

Electronic Spectra and Electrochemical Data

Complex	$\lambda_{\max}$ , nm ( $\log \epsilon$ ) <sup>a</sup>	$E_{p/2}$ (V) vs sce <sup>b</sup>
I	725(1.60), 545(3.30), 418(3.36), 349(3.64), 275(sh, 4.20), 237(4.46), 206(4.35)	-0.59 red +1.36 ox <sup>c</sup>
II	790(1.60), 594(3.445), 447(3.425), 371(3.72), 280(sh, 4.32), 248(4.415), 211(4.41)	-0.70 red +1.07 ox <sup>c</sup> +1.76 ox <sup>c</sup>
III	756(1.70), 575(sh, 2.34), 419(3.16), 350(sh, 3.60), 248(4.505), 223(4.90)	d

<sup>a</sup> measured in spectroquality CH<sub>3</sub>CN.


<sup>b</sup> cyclic voltammetry in CH<sub>3</sub>CN with 0.1 M (C<sub>2</sub>H<sub>5</sub>)<sub>4</sub>N<sup>+</sup>PF<sub>6</sub><sup>-</sup> supporting electrolyte, platinum button electrode; reversible waves except where noted; red = reduction, ox - oxidation.

<sup>c</sup> irreversible wave.

<sup>d</sup> all waves irreversible, none well-defined.



Figure 1. The proposed structure and 80.5 MHz  $^{11}\text{B}$  nmr spectrum  
of I, 4,5-( $\eta^5\text{-C}_5\text{H}_5$ ) $_2$ -4,5- $\text{Fe}_2$ -2,3- $\text{C}_2\text{B}_9\text{H}_{11}$ .



● CH  
○ BH

-113.1

(1)

-42.4

(2)

-22.4

(4)

-5.7

(2)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report 98	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Synthesis of Electron-Deficient Biferracarboranes		5. TYPE OF REPORT & PERIOD COVERED Interim
7. AUTHOR(s) C. G. Salentine and M. F. Hawthorne		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS The University of California Department of Chemistry Los Angeles, California 90024		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0390
11. CONTROLLING OFFICE NAME AND ADDRESS Chemistry Branch Office of Naval Research Washington, D.C. 20360		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-608
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE January 17, 1978
		13. NUMBER OF PAGES 14 (incl. fig.)
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Electron-deficient Biferracarboranes Diamagnetic Ferric Long-range coupling		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The polyhedral expansion of $3,1,2-C_5H_5FeC_2B_9H_{11}$ produced three new electron-deficient biferracarboranes $(C_5H_5Fe)_2C_2B_9H_{11}$ , $(C_5H_5Fe)_2C_2B_9H_9(OH)$ and $[C_5H_5Fe-C_2B_9H_{11}FeC_2B_9H_{11}]^-$ . All three complexes contained formal Fe(III) atoms and were diamagnetic. The proposed structures, based on spectroscopic and crystallographic data, contain non-bonded iron atoms. These complexes constitute the first examples of long-range electron-spin coupling through a carborane polyhedron.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)



APPENDIX



# TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
Office of Naval Research Arlington, Virginia 22217 Attn: Code 102IP 1	6	U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. Jerry Smith	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Civil Engineering Laboratory Port Hueneme, California 93041 Attn: Mr. W. S. Haynes	1
ONR Branch Office 760 Market St., Rm. 447 San Francisco, Calif. 94102 Attn: Dr. P. A. Miller	1	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
ONR Branch Office 495 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Director Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Office of Naval Research Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
The Asst. Secretary of the Navy (R&D) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Dr. R. M. Grimes University of Virginia Department of Chemistry Charlottesville, Virginia 22901	1
Commander Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1	Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1
		Dr. C. Quicksall Georgetown University Department of Chemistry 37th & O Streets Washington, D.C. 20007	1

TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1
Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916	1		
Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712	1		
Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1		
Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1		
Dr. M. H. Chisholm Princeton University Department of Chemistry Princeton, New Jersey 08540	1		
Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1		
Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201	1		